

Title: Progenitor cell markers predict outcome of patients with Hepatocellular Carcinoma beyond Milan criteria undergoing liver transplantation

Authors: Oriana Miltiadous¹, Daniela Sia^{1,2}, Yujin Hoshida¹, Maria Isabel Fiel¹, Andrew N. Harrington¹, Swan N. Thung¹, Poh Seng Tan^{1,3}, Hui Dong⁴, Kate Revill¹, Charissa Y Chang¹, Sasan Roayaie⁵, Thomas J Byrne⁶, Vincenzo Mazzaferro², Jorge Rakela⁶, Sander Florman¹, Myron Schwartz¹, and Josep M Llovet^{1,7,8}.

Affiliation:

¹ Mount Sinai Liver Cancer Program (Division of Liver Diseases, Department of Medicine, Tisch Cancer Institute, Department of Pathology, Recanati Miller Transplantation Institute, Department of Surgical Oncology), Icahn School of Medicine at Mount Sinai, New York, USA.

² Gastrointestinal Surgery and Liver Transplantation Unit, Department of Surgery, National Cancer Institute, Milan, Italy.

³ Division of Gastroenterology and Hepatology, University Medicine Cluster, National University Health System, Singapore

⁴ Eastern Hepatobiliary Surgery Hospital, Second Military Medical University Shanghai, China

⁵ Liver Cancer Program, Hofstra-North Shore LIJ School of Medicine, Lenox Hill Hospital, New York, USA

⁶ Division of Hepatology, Mayo Clinic, Phoenix, AZ, USA.

⁷ Liver Cancer Translational Research Laboratory, Barcelona - Clínic Liver Cancer Group (Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Liver Unit. Hospital Clínic, Universitat de Barcelona, Catalonia, Spain.

⁸ Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain.

Address for correspondence:

Josep M Llovet, MD
Mount Sinai Liver Cancer Program
Division of Liver Diseases,
Tisch Cancer Institute
Icahn School of Medicine at Mount Sinai.
Madison Ave 1425. 11F-70. Box 1123
New York, NY10029. USA
Phone: 212-6599503
FAX: 212-849-2574
E-mail: Josep.Llovet@mssm.edu

Electronic word count

Characters count Title: 121

Words count Abstract: 251

Words count includes Abstract, Introduction, Materials and Methods, Results,
Discussion, References, Tables and Figure Legends: 5985
References: 45

Number of Figures and Tables: 3 Figures and 5 Tables

Supplementary Material: 6 Figures and 6 Tables

List of abbreviations

HCC: Hepatocellular carcinoma
LT: Liver transplantation

1 CK19: Cytokeratin protein 19

2
3 HR: Hazard Ratio:

4
5 EpCAM: Epithelial cell adhesion molecule

6
7
8 OS: Overall survival

9
10 miVI: microvascular invasion

11
12 TGFb: Transforming growth factor beta

13
14 CT: Computed Tomography

15
16
17 MRI: Magnetic Resonance Imaging

18
19
20 AFP: Alpha-fetoprotein

21
22 IHC, immunohistochemical

23
24 RPS6: ribosomal protein S6

25
26
27 FDR: False discovery rate

28
29
30 H&E: Hematoxylin and eosin

31
32 PPV: Positive predictive value

33
34
35 NPV: Negative predictive value

36
37 HCV: Hepatitis C virus

38
39
40 HBV: Hepatitis B Virus

41
42 VIF: Variance Inflation Factor

43
44
45
46
47 **Keywords:** gene expression, prognosis, stem cell, gene signature, survival

48
49
50
51
52 **GEO Accession number:** GSE62743

53
54
55
56
57 **Conflict of Interest:** the authors indicate no potential conflicts of interest.

Financial Support:

Josep M. Llovet was supported by grants from The European Commission Framework Programme 7 (HEPTROMIC, Proposal No: 259744), the Samuel Waxman Cancer Research Foundation, the Spanish National Health Institute (SAF2013-41027), and the Asociación Española Contra el Cáncer (AECC). Daniela Sia and Vincenzo Mazzaferro are supported by the Italian Association for Cancer Research and the Italian National Ministry of Health. Yujin Hoshida is supported by grants from the U.S. National Institute of Diabetes and Digestive and Kidney Diseases (R01 DK099558), the European Commission Framework Programme 7 (HEPTROMIC, Proposal No: 259744).

Author's contributions

Conceived and designed the experiments: MS, JML, and JR

Performed the experiments: OM, DS, YH, ANH, PST, HD, and KR

Analyzed the data: OM, DS, YH, MIF, SNT, PST, CYC, JML

Contributed reagents/materials/analysis tools: OM, YH, ANH, PST, CYC, SR, TJB, VM, JR, SF, MS

Wrote the manuscript: OM, DS, YH, MS, JML

Clinical data: OM, ANH, CHC, SR, VM, JR, SF, MS, JML

Obtained funding: JR, SF, MS, JML

Critical review of the manuscript: OM, DS, YH, MIF, ANH, SNT, PST, KR, SR, TJB, VM, JR, SF, MS and JML

ABSTRACT

Background and Aims: In patients with hepatocellular carcinoma (HCC), liver transplantation (LT) is an excellent therapy if tumor characteristics are within the Milan criteria. We aimed to define genomic features enabling to identify HCC patients beyond Milan criteria who have acceptable transplant outcomes.

Methods: Among 770 consecutive HCC patients transplanted between 1990 and 2013, 132 had tumors exceeding Milan criteria on pathology and were enrolled in the study; 44% of the patients satisfied the 'up-to-7 rule' [7=sum of the size of the largest tumor and the number of tumors]. Explant tumors were assessed for genomic signatures and immunohistochemical markers associated with poor outcome.

Results: At a median follow-up of 88 months, 64 patients had died and 45 recurred; the 5-year overall survival (OS) and recurrence rates were 57% and 35%, respectively. Cytokeratin 19 (CK19) gene signature was independently associated with recurrence [Hazard ratio (HR)=2.95, $p<0.001$], along with tumor size (HR=3.37, $p=0.023$) and presence of satellites (HR=2.98, $p=0.001$). S2 subclass signature was independently associated with poor OS (HR=3.18, $p=0.001$), along with tumor size (HR=5.06, $p<0.001$) and up-to-7 rule (HR=2.50, $p=0.002$). Using the presence of progenitor cells markers (either CK19 or S2 signatures) patients were classified into poor-prognosis (n=58; 5-year recurrence 53%, survival 45%) and good-prognosis (n=74; 5-year recurrence 19%, survival 67%) (HR=3.16, $p<0.001$ for recurrence, and HR=1.72, $p=0.04$ for OS).

Conclusions:

HCC patients transplanted beyond Milan criteria without gene signatures of progenitor markers (CK19 and S2) achieved survival rates similar as those within Milan criteria.

Once prospectively validated, these markers may support a limited expansion of LT indications.

INTRODUCTION

Liver transplantation (LT) is an effective treatment option for hepatocellular carcinoma (HCC) when disease is defined by the widely accepted Milan criteria [1, 2]. Transplantation for patients within the Milan criteria generally yields a 5-year overall survival (OS) of 70% and a recurrence rate less than 15% [1-3]. Several efforts have been made to expand the criteria based on tumor size and number [4-8]. Although downstaging of tumors beyond Milan criteria is accepted by some UNOS regions, it has not yet been adopted by international consensus guidelines of LT [9] or guidelines of management of HCC [2]. Mazzaferro et al. proposed the up-to-7 rule [the sum of the number of tumor nodule(s) and the maximum diameter of the nodule(s) must not exceed the value of 7] which, in the absence of microvascular invasion (miVI) results in 5-year OS above 70% [8]. This study along with previously published data confirmed miVI to be a key predictor of recurrence in patients with HCC [10]. However, this information cannot be used in pre-transplant decision-making as an indication of transplantation since miVI is only diagnosed based on post-surgical histological assessment. This highlights the limitations of the current image-based prognostic algorithm for selecting HCC patients for LT.

Genome-wide transcriptome profiling has identified several key deregulated genes, molecular pathways and signatures associated with disease progression and prognosis in HCC [11, 12]. Activation of specific molecular pathways such as transforming growth factor-beta (TGFB) as well as presence of progenitor cell markers such as cytokeratin 19 (CK19) and epithelial cell adhesion molecule (EPCAM) have been associated with more aggressive biological tumor characteristics and rapid disease spread [13-35].

1 In a previous study, we identified gene signatures that significantly improved prediction
2 of HCC recurrence after surgical resection [21]. Here we sought to define if gene
3 signatures are also able to identify patients with HCC beyond Milan criteria who
4 nevertheless may have acceptable outcomes with LT. In addition, we are providing the
5 transcriptomic landscape of patients at more advanced states of the disease
6 compared with those undergoing resection or transplantation according to Milan
7 criteria, as per AASLD/EASL guidelines [2, 36].
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

MATERIALS AND METHODS

Patient cohorts and tissue samples

A total of 770 patients were transplanted for HCC between 1990 and 2013 at Mount Sinai Hospital, New York (n=590) and Mayo Clinic, Scottsdale-Arizona (n=180). Among these, we selected 132 patients (Mount Sinai: n=94, Mayo: n=38) with HCC beyond Milan criteria by pathological assessment, for whom base-line and follow-up clinical information and archived fixed tissue of viable tumor were available (**Figure 1**). The study protocol was approved by the respective Institutional Review Boards. See Supplementary Methods.

Genome-wide transcriptome profiling

Total RNA was subjected to transcriptome profiling using Whole-Genome DASL-HT (Illumina). See Supplementary Methods. GEO accession number: GSE62743.

Immunohistochemistry

Immunohistochemistry was performed for the evaluation of nuclear β -catenin as a surrogate for Wnt pathway activation; phosphorylation of ribosomal protein S6 (RPS6) as the major downstream effector of activated AKT-mTOR pathway; CK19 and EPCAM as surrogate of progenitor cell origin. See Supplementary Methods.

Bioinformatics and statistical analysis

Presence of previously reported HCC gene signatures (**Supplementary Table 1**) was evaluated (**Figure 2**). See Supplementary Methods.

RESULTS

Patient characteristics

Clinico-pathological data and outcomes of the patients are summarized in **Table 1**. Most of the patients were male (80%) and hepatitis C virus (HCV)-infected (71%). The median size of the largest tumor was 4 cm (range 1-9 cm). Tumors exceeded Milan criteria due to single tumor larger than 5 cm (5%), 2 to 3 tumors with the largest tumor larger than 3 cm (39%), or more than 3 tumors within the liver (57%). Satellite lesions (defined as HCC nodules < 2 cm in diameter within 2 cm from main tumor [37]) were present in 18% and miVI was present in 84% of the patients. Serum alpha fetoprotein (AFP) was > 200 mg/dL in 17% of cases. At the end of follow-up, 64 patients had died, OS at 5- and 10-years was 57% and 42%, respectively. HCC recurrence developed in 45 patients, 5 and 10-year cumulative recurrence rates were 35% and 44%, respectively (**Figure 3**).

Clinical variables identified as predictors of recurrence on multivariate analysis included tumor diameter >5 cm [$p=0.023$, Hazard Ratio (HR) = 3.37] and the presence of satellites ($p<0.001$, HR=2.98). Tumor diameter >5 cm ($p<0.001$, HR=5.06) and outside up-to-7 rule (HR=2.50, $p=0.002$) predicted poor survival (**Tables 2 and 3**).

Prevalence of prognostic liver cancer gene signatures

Among the 22 liver cancer gene signatures tested, 16 were able to sort patients into good and poor prognosis groups with a FDR<0.05 (**Figure 2A**). The human [21] and rat [28] CK19 gene signatures were present in 40% and 42% of patients, respectively. Signatures associated with poor clinical outcome and/or more aggressive biological features were generally present in the same set of tumors. At least one poor-prognosis signature was present in 84 patients (64%), this being a higher rate than we have

previously reported in patients treated with resection [21]. Twenty-five tumors (19%) harbored more than 10 poor prognosis/aggressiveness gene signatures. VI signature was positive in 22% of cases, 93% of which were positive for histological miVI. The majority of tumors lacking poor prognosis/aggressive signatures were enriched by the S3 subclass signature (39/48, 81%), which is associated with smaller, higher proportion of well-differentiated tumors [13].

The gene signatures analyzed were consistent with 3 distinct clusters or major molecular classes of HCC (**Figure 2B**): (1) *Proliferation and cell cycle activation* (Cluster A [16], Cholangiocarcinoma [29], Proliferation [20], Recurrence [24], S1 [13], TGFb [23], MET [18], and G3 [19] signatures), 2) *Proliferation-progenitor cell origin* (Hepatoblastoma C2 [26], Class S2 [13], EpCAM [25] and VI [22] signatures), which includes a subclass of CK19 positive tumors (CK19 human [21] and rat [28] signatures), and 3) *non-proliferative class*, enriched by tumors positive for CTNNB1[20] or class S3 [13, 33]. Interestingly, there was no association between the underlying etiology [HCV, Hepatitis B Virus (HBV) or alcohol] and the expression of signatures.

Comparing the classification of signatures between this cohort and previously published results from resection cases [21], we noticed a higher frequency of tumors positive for ≥ 6 signatures of poor prognosis (32.6% vs 25%, respectively). Similarly, our cohort presented a higher prevalence of CK19 human signatures (41% vs 34%, respectively), suggesting that it includes more advanced tumors compared with our previous report on resected patients [21].

Immunohistochemical (IHC) analysis

CK19 immunostaining was positive in 6 tumors (5%), and was significantly associated with the presence of the human ($p=0.03$) and rat ($p=0.04$) CK19 gene signatures, as well as VI signature [22] ($p<0.01$) (**Supplementary Table 2**). EPCAM immunostaining was positive in 20 tumors (15%), and was significantly associated with EPCAM signature ($p=0.01$), human and rat CK19 signatures ($p=0.01$ and $p=0.02$, respectively), Proliferation subclass signature ($p=0.01$), and VI signature ($p=0.01$) (**Supplementary Table 2**). Nuclear staining of β -catenin was observed in 27 tumors (21%), and was positively associated with CTNNB1 class signature ($p=0.01$) (**Supplementary Figure 1, Supplementary Table 2**). In addition, 12 samples (9%) had overexpression of cytoplasmic β catenin with one third of these in CTNNB1 class (4/12, 33%). Phospho-RPS6 staining was positive in 75 tumors (57%) with no obvious correlation with any of the evaluated gene signatures.

Prognostic relevance of liver cancer gene signatures in LT treated patients beyond Milan

Prognostic gene signatures in HCC were initially developed in patients undergoing resection, a treatment indicated for early stages of the disease. Univariate Cox regression analysis demonstrated that VI [22], G3 subclass [19], and S2 subclass [13] signatures were associated with overall survival, along with patient's age, tumor size, and up-to-7 rule (**Table 2**). In the multivariate analysis only S2 signature [13] ($HR=3.18$, $p=0.001$), tumor diameter >5 cm ($HR=5.06$, $p<0.001$), and outside up-to-7 rule ($HR=2.50$, $p=0.002$) were independent predictors of survival. VI [22], G3 [19], Proliferation [20], CK19 human [21], CK19 rat [28], Cholangiocarcinoma [29], Cluster A [16] and S3 [13] gene signatures were associated with HCC recurrence in univariate

analysis along with tumor size, AFP and satellites from clinical variables as well as RPS6 from IHC (**Table 3**). In multivariate analysis, CK19 human [21] (HR=2.91, $p=0.001$), tumor diameter > 5 cm (HR=3.37, $p=0.023$) and satellites (HR=2.98, $p<0.001$) were independent predictors of recurrence.

Of note, histological miVI predicted neither survival nor recurrence, presumably because in this cohort it was present in the large majority of cases (112 patients, 84.8%). Also, IHC positivity for reported progenitor cell markers also failed to improve outcome prediction.

Combination of progenitor cell signatures (CK19 and S2) discriminates prognostic subgroups in transplant patients

Multivariate analyses demonstrated prognostic significance of two progenitor-related gene signatures, CK19 [21] and S2 [13]. Furthermore, S2 subclass signature was also able to predict significant shorter survival post-recurrence in a subgroup analysis, suggesting that this signature might be able to capture more aggressive tumors with high risk of recurrence ($p<0.001$, HR= 5.11, **Supplementary Figure 2**). We therefore explored whether combining these signatures could further improve our ability to predict outcome. At least one of these two signatures was present in 58 patients (44%), and having one or both signatures independently predicted both OS (HR=1.72, $p=0.036$) and recurrence (HR=3.16, $p=0.001$) (**Tables 4 and 5**): patients with one or both signatures had 45% 5-year OS and 53% recurrence vs 67% 5-year OS and 19% recurrence when neither signature was present (**Figure 3C-D**). The presence of any of the two signatures also predicted recurrence when the analysis was limited to patients beyond Milan criteria based on preoperative imaging ($n=94$) (**Supplementary Figure 3, Supplementary Table 3**). Overall, 5-year recurrence rate was of 35.7% and was

significantly lower in patients CK19/S2 rule negative (20.8% vs 52.8%, $p=0.003$).
Furthermore, the CK19/S2 rule, when applied to patients classified according to BCLC stage, significantly improved the prognostic prediction by stratifying patients in high and low risk for both recurrence and survival (**Supplementary Figures 4 and 5**).

Discussion

Almost 20 years after the seminal observation by Mazzaferro et al [1], we continue to rely on size and number of tumors for selecting patients with HCC for LT. Since then, the concept of a modest expansion beyond the Milan criteria has been proposed without gaining global acceptance [4, 38-39]. Most of the studies performed were retrospective, used pathological data and only included few patients with HCC beyond Milan criteria, thus leading to unrealistic good results because the outcome was “diluted” among a much larger pool of patients within Milan criteria. The largest series ever reported exploring expansion of Milan criteria led to the pathological up-to-7 rule where outcomes were significantly better for patients within this rule, as long as vascular invasion is not present [8]. After all, tumor size and number remain surrogates for the biological aggressiveness of HCC; henceforth more refined markers are needed.

While there have been multiple publications linking various genetic profiles of HCC with outcomes after hepatic resection, tumor biology in transplanted patients has not been assessed widely thus far. Most studies have relied on degree of HCC differentiation on biopsy, AFP and response to bridge therapy while on the waiting list [40-42]. Using allelic imbalance, we described that genomic instability of certain microsatellites selected for proximity to known oncogenes can help select patients beyond Milan criteria who are likely to have favorable outcomes after transplantation [43]. Because gene expression profiling could be eventually performed on needle liver biopsies in a pre-transplant setting [44], the positivity for each of the signatures can assist in the decision making of selecting patients for transplantation.

In our current study, we explored the tumor gene expression profiles for the presence of 22 previously reported prognostic signatures –reflecting HCC molecular

subclasses or activation of signaling pathways- in 132 patients transplanted with HCC beyond Milan criteria and key immunohistochemical staining and correlated the results with clinical variables and outcome. Sixteen signatures demonstrated association with recurrence and/or survival; among these the CK19 and the S2 signatures, both denoting progenitor cell features, were independent predictors of outcome when included in models incorporating clinical variables. As shown in **Figure 2**, both signatures cluster together along with others from hepatoblastoma, which overall reflect progenitor cell origin. In fact, CK19 is a molecular marker of progenitor cells, whereas S2 subclass [13] has been linked with high levels of expression of AFP even at early stages of the disease, and activation of mTOR and IGF signaling [21, 35]. We also found enrichment for Met and TGFb signaling in these patients. In our study, one and/or both signatures were present in 44% of cases and the overall 5 yr-survival and recurrence rate (53% and 45%, respectively) was discouraging. Conversely, in the 56% of patients harboring neither signature, OS was 67% and recurrence was 19% at 5 years, similar to the results of LT for HCC when conventional Milan criteria are applied. The capacity of our tool to rule out tumors with “poor biology” is directly reflected in the acceptable low recurrence rate.

In order to be used prospectively, the CK19/S2 gene signature needs to be validated when Milan criteria are defined on pre-operative imaging. Indeed, in the subgroup analysis of pre-operatively defined cases beyond Milan criteria, the CK19/S2 signature remained independently associated with recurrence. However, the signature no longer retained significance on multivariate analysis for overall survival, likely due to the smaller sample size. Overall, these findings support the possibility that genomic analysis of HCC performed prior to LT may enable expansion of the current imaging-

1 based selection criteria to include patients beyond Milan Criteria but with favorable
2 genomic profiles, without significantly impacting clinical outcome.
3

4
5 Another interesting finding of our study was the relative genomic disparity observed in
6
7 HCC cases treated with LT versus those undergoing resection. In this series of
8
9 patients with HCC beyond Milan Criteria we observed a higher frequency of tumors
10
11 demonstrating ≥ 6 signatures than we did in the study of patients undergoing liver
12
13 resection [21]. This may be explained, to some degree, by the more advanced clinical
14
15 stage of cancers that received LT beyond Milan criteria versus those undergoing
16
17 resection. While resection is limited to single tumors, nearly 95% of the patients in the
18
19 current study were transplanted for multiple HCC.
20
21
22
23

24
25 We intend to validate these observations prospectively, but in order to establish
26
27 the absence of the CK19 and S2 progenitor cell signatures as a basis for safe
28
29 expansion of the indication for LT for patients with HCC, a number of barriers must be
30
31 overcome. First, the study will have to be limited to patients identified pre-LT as being
32
33 beyond Milan Criteria in order to be clinically relevant. The genomic heterogeneity of
34
35 tumors is widely recognized, all the more so when the tumor is large or multinodular as
36
37 in patients beyond Milan Criteria. While we have shown in a limited study that genomic
38
39 heterogeneity in HCC rarely results in assignment of samples from the same tumor to
40
41 different genomic classes [21], the degree to which a pre-LT biopsy accurately reflects
42
43 the genomic landscape of the tumor remains unclear. Molecular heterogeneity
44
45 between and within tumors can pose a risk for miss-classifications when selecting the
46
47 area to biopsy prior LT. However, with the advancements of imaging techniques, a
48
49 significant correlation has been shown between tumor enhancement and histological
50
51 differentiation which could greatly assist in more accurate profiling [45].
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 On the other side, the impact on gene expression of pre-LT nonsurgical tumor
2 treatment, that is commonly performed to down-stage HCC and forestalls progression
3 while awaiting transplant, remains undefined. Furthermore, with long pre-LT waiting
4 times before transplantation and the potential drop out of some patients, the stability of
5 tumor gene expression profiles over time must be considered. Thus, the extent to
6 which a biopsy done today reflects the nature of the tumor later on and the frequency
7 with which biopsy will need to be repeated, need to be determined.

8
9
10
11
12
13
14
15
16
17 Despite its shortcomings, our observations provide initial insights into the
18 understanding of tumor biology and clinical outcomes beyond size and number of
19 HCC nodules. Those patients with tumors with “good biology” as defined by absence
20 of signatures with progenitor cell traits (CK19/S2) represent half of the patients
21 transplanted beyond Milan criteria in 2 institutions in the United States. Our seminal
22 study provides the rationale for expanding the conventional criteria based upon
23 genomic data, since outcomes are competitive with those achieved by HCC patients
24 within Milan criteria. It is a novel, genomic, complementary tool to refine previously
25 reported proposals, such as Up-to-7 rule [8]. Certainly, these results need to be
26 validated using a separate external cohort of patients and then examined
27 prospectively before they can be adopted by clinical guidelines.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES

- [1] Mazzaferro V, Regalia E, Doci R, Andreola S, Pulvirenti A, Bozzetti F et al. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med* 1996;334:693-700.
- [2] Llovet JM, Ducreux M, Lencioni R, Di Bisceglie AM, Galle PR, Dufour JF et al. J EASL-EORTC clinical practice guidelines: management of hepatocellular carcinoma. *Hepatology* 2012;56:908-43.
- [3] Llovet JM, Fuster J, Bruix J. Intention-to-treat analysis of surgical treatment for early hepatocellular carcinoma: resection versus transplantation. *Hepatology* 1999;30:1434-40.
- [4] Yao FY, Ferrell L, Bass NM, Watson JJ, Bacchetti P, Venook A, et al. Liver transplantation for hepatocellular carcinoma: expansion of the tumor size limits does not adversely impact survival. *Hepatology* 2001;33:1394-403.
- [5] Roayaie S, Frischer JS, Emre SH, Fishbein TM, Sheiner PA, Sung M et al. Long-term results with multimodal adjuvant therapy and liver transplantation for the treatment of hepatocellular carcinoma larger than 5 centimeters. *Ann Surg.* 2002;235:533–539.
- [6] Lee SG, Hwang S, Moon DB, Ahn CS, Kim KH, Sung KB et al. Expanded Indication Criteria of Living Donor Liver transplantation for Hepatocellular Carcinoma at One Large-Volume Center. *Liver Transpl.* 2008;14:935-45.
- [7] Yao FY. Liver transplantation for hepatocellular carcinoma: Beyond the Milan criteria. *Am J Transplant* 2008;8:1982-9.
- [8] Mazzaferro V, Llovet JM, Miceli R, Bhoori S, Schiavo M, Mariani L et al. Predicting survival after liver transplantation in patients with hepatocellular carcinoma beyond the Milan criteria: a retrospective, exploratory analysis. *Lancet Oncol.* 2009;10:35-43.

- [9] Clavien PA, Lesurtel M, Bossuyt PM, Gores GJ, Langer B, Perrier A; OLT for HCC Consensus Group. Recommendations for liver transplantation for hepatocellular carcinoma: an international consensus conference report. *Lancet Oncol* 2012;13:e11-22.
- [10] Jonas S, Bechstein WO, Steinmüller T, Herrmann M, Radke C, Berg T, et al. Vascular invasion and histopathologic grading determine outcome after liver transplantation for hepatocellular carcinoma in cirrhosis. *Hepatology* 2001;33:1080-6.
- [11] Villanueva A, Hernandez-Gea V, Llovet JM. Medical therapies for hepatocellular carcinoma: a critical view of the evidence. *Nature Rev Gastroenterol Hepatol* 2013;10:34-42.
- [12] Sawyer CL. The cancer biomarker problem, *Nature* 2008;452:548-52.
- [13] Hoshida Y, Nijman SM, Kobayashi M, Chan JA, Brunet JP, Chiang DY et al. Integrative Transcriptome Analysis Reveals Common Molecular Subclasses of Human Hepatocellular Carcinoma. *Cancer Res* 2009;69:7385-92.
- [14] Ye QH, Qin LX, Forgues M, He P, Kim JW, Peng AC et al. Predicting hepatitis B virus-positive metastatic hepatocellular carcinomas using gene expression profiling and supervised machine learning. *Nat Med* 2003;9:416-23.
- [15] Iizuka N, Oka M, Yamada-Okabe H, Nishida M, Maeda Y, Mori N et al. Oligonucleotide microarray for prediction of early intrahepatic recurrence of hepatocellular carcinoma after curative resection. *Lancet* 2003;361:923-9.
- [16] Lee JS, Chu IS, Heo J, Calvisi DF, Sun Z, Roskams T et al. Classification and prediction of survival in hepatocellular carcinoma by gene expression profiling. *Hepatology* 2004;40:667-76.

- [17] Kurokawa Y, Matoba R, Takemasa I, Nagano H, Dono K, Nakamori S et al. Molecular-based prediction of early recurrence in hepatocellular carcinoma. *J Hepatol*. 2004;41:284-91.
- [18] Kaposi-Novak P, Lee JS, Gómez-Quiroz L, Coulouarn C, Factor VM, Thorgeirsson SS. Met-regulated expression signature defines a subset of human hepatocellular carcinomas with poor prognosis and aggressive phenotype. *J Clin Invest*, 2006;116:1582-95.
- [19] Boyault S, Rickman DS, de Reyniès A, Balabaud C, Rebouissou S, Jeannot E, et al. Transcriptome classification of HCC is related to gene alterations and to new therapeutic targets. *Hepatology* 2007;45:42-52.
- [20] Chiang DY, Villanueva A, Hoshida Y, Peix J, Newell P, Minguez B, et al. Focal gains of VEGFA and molecular classification of hepatocellular carcinoma. *Cancer Res*. 2008;68:6779-88.
- [21] Villanueva A, Hoshida Y, Battiston C, Tovar V, Sia D, Alsinet C, et al. Combining clinical, pathology, and gene expression data to predict recurrence of hepatocellular carcinoma. *Gastroenterology*, 2011;140:1501-12.
- [22] Mínguez B, Hoshida Y, Villanueva A, Toffanin S, Cabellos L, Thung S et al. Gene-expression signature of vascular invasion in hepatocellular carcinoma. *J Hepatol*. 2011;55:1325-31.
- [23] Coulouarn C, Factor VM, Thorgeirsson SS. Transforming growth factor-beta gene expression signature in mouse hepatocytes predicts clinical outcome in human cancer. *Hepatology* 2008;47:2059-67.
- [24] Woo HG, Park ES, Cheon JH, Kim JH, Lee JS, Park BJ, et al. Gene expression-based recurrence prediction of hepatitis B virus-related human hepatocellular carcinoma. *Clin Cancer Res* 2008;14:2056-64.

- [25] Yamashita T, Forgues M, Wang W, Kim JW, Ye Q, Jia H, et al. EpCAM and alpha-fetoprotein expression defines novel prognostic subtypes of hepatocellular carcinoma. *Cancer Res* 2008;68:1451-61.
- [26] Cairo S, Armengol C, De Reyniès A, Wei Y, Thomas E, Renard CA et al. Hepatic stem-like phenotype and interplay of Wnt/beta-catenin and Myc signaling in aggressive childhood liver cancer. *Cancer Cell*, 2008;14:471-84.
- [27] Yoshioka S, Takemasa I, Nagano H, Kittaka N, Noda T, Wada H et al. Molecular prediction of early recurrence after resection of hepatocellular carcinoma. *Eur J Cancer* 2009;45:881-9.
- [28] Andersen JB, Loi R, Perra A, Factor VM, Ledda-Columbano GM, Columbano A et al. Progenitor-derived hepatocellular carcinoma model in the rat. *Hepatology* 2010;51:1401-9.
- [29] Woo HG, Lee JH, Yoon JH, Kim CY, Lee HS, Jang JJ et al. Identification of a cholangiocarcinoma-like gene expression trait in hepatocellular carcinoma. *Cancer Res.* 2010;70:3034-41.
- [30] Roessler S, Long EL, Budhu A, Chen Y, Zhao X, Ji J et al. Integrative genomic identification of genes on 8p associated with hepatocellular carcinoma progression and patient survival. *Gastroenterology* 2012;142:957-966.
- [31] Nault JC, De Reyniès A, Villanueva A, Calderaro J, Rebouissou S, Couchy G et al. A hepatocellular carcinoma 5-gene score associated with survival of patients after liver resection. *Gastroenterology* 2013;145:176-87.
- [32] Komuta M, Spee B, Vander Borgh S, De Vos R, Verslype C, Aerts R et al. Clinicopathological study on cholangiolocellular carcinoma suggesting hepatic progenitor cell origin. *Hepatology* 2008;47:1544-56.

- [33] Lachenmayer A, Alsinet C, Savic R, Cabellos L, Toffanin S, Hoshida Y et al. Wnt-pathway activation in two molecular classes of hepatocellular carcinoma and experimental modulation by sorafenib. *Cancer Res* 2012;18:4997-5007
- [34] Bjornsti MA, Houghton PJ. The tor pathway: a target for cancer therapy. *Nat Rev Cancer*;4:335-48.
- [35] Villanueva A, Chiang DY, Newell P, Peix J, Thung S, Alsinet C et al. Pivotal role of mTOR signaling in hepatocellular carcinoma. *Gastroenterology* 2008;135:1972-83.
- [36] Bruix J, Sherman M; American Association for the Study of Liver Diseases. Management of hepatocellular carcinoma: an update. *Hepatology*. 2011;53:1020-2.
- [37] Roayaie S, Blume IN, Thung SN, Guido M, Fiel MI, Hiotis S et al. A system of classifying microvascular invasion to predict outcome after resection in patients with hepatocellular carcinoma. *Gastroenterology* 2009;137:850-5.
- [38] Kim PT, Onaca N, Chinnakotla S, Davis GL, Jennings LW, McKenna GJ, et al. Tumor biology and pre-transplant locoregional treatments determine outcomes in patients with T3 hepatocellular carcinoma undergoing liver transplantation. *Clin Transplant*. 2013;27:311-8
- [39] Cillo U, Vitale A, Grigoletto F, Gringeri E, D'Amico F, Valmasoni M, et al. Intention-to-treat analysis of liver transplantation in selected, aggressively treated HCC patients exceeding the Milan criteria. *Am J Transplant*. 2007;7:972-81.
- [40] DuBay D, Sandroussi C, Sandhu L, Cleary S, Guba M, Cattral MS, et al. Liver transplantation for advanced hepatocellular carcinoma using poor tumor differentiation on biopsy as an exclusion criterion. *Ann Surg*. 2011;253:166-72.
- [41] Toso C, Asthana S, Bigam DL, Shapiro AM, Kneteman NM. Reassessing selection criteria prior to liver transplantation for hepatocellular carcinoma utilizing the Scientific Registry of Transplant Recipients database. *Hepatology* 2009;49:832-8

- 1 [42] Yao FY, Kerlan RK Jr, Hirose R, Davern TJ, Bass NM, Feng S, et al. Excellent
2 outcome following down-staging of hepatocellular carcinoma prior to liver
3 transplantation: an intention-to-treat analysis. *Hepatology*. 2008;48:819-27.
4
5 [43] Schwartz M, Dvorchik I, Roayaie S, Fiel MI, Finkelstein S, Marsh JW, et al. Liver
6 transplantation for hepatocellular carcinoma: extension of indications based on
7 molecular markers. *J Hepatol*. 2008;49:581-8.
8
9 [44] Hoshida Y, Villanueva A, Kobayashi M, Peix J, Chiang DY, Camargo A et al.
10 Gene expression in fixed tissues and outcome in hepatocellular carcinoma. *N Engl J*
11 *Med* 2008;359:1995-2004.
12
13 [45] Banerjee S, Wang DS, Kim HJ, Sirlin CB, Chan MG, Korn RL et al. A computed
14 tomography radiogenomic biomarker predicts microvascular invasion and clinical
15 outcomes in hepatocellular carcinoma. *Hepatology* 2015 Apr 30.

Acknowledgements

We thank Laia Cabellos for technical assistance and Xiaochen Sun for bioinformatic support.

FIGURE LEGENDS

Figure 1: Flow chart of the study. The initial cohort included 770 patients that were consecutively transplanted for HCC at Mount Sinai Hospital, New York (n=590) and Mayo Clinic, Scottsdale Arizona (n=180). A total of 206 patients were beyond Milan criteria based on the explant pathology. Tumors with HCC-ICC features, diffused pattern, necrotic tissue, macro-vascular invasion or metastasis were excluded. From the remaining 141 cases, 6 tumors with poor-quality RNA were discarded. Total RNA from 135 tumors was subjected to transcriptome profiling. Three samples had poor quality profile and were excluded and eventually 132 tumors were tested for the presence of previously reported outcome-associated gene signatures.

Figure 2: Prediction of gene expression signatures. Only signatures that were able to assign patients into good and poor prognosis groups with $FDR > 0.05$ were included. **(A)** Each column represents different sample and each row different signature. Positivity of each signature is represented by red bars. Events (recurrence or 5-year death) are shown with black bars. **(B)** Visualization of cramer's V coefficient for the pair-wise comparison of gene signatures. The scale from blue to red represents the strength of correlation (red represents the highest correlation). The signatures are clustered according to their correlation.

Figure 3: Kaplan-Meier curves and estimates of overall survival and recurrence in patients beyond Milan undergoing liver transplantation (n=132). Top figures show overall survival and recurrence in this cohort. Patients positive for either of CK19 or S2 signatures (progenitor-cell signatures) were assigned in a poor-outcome group.

Lower panels show overall survival and recurrence in patients beyond Milan with or without CK19/S2 signatures. P-values are calculated based on the log-rank test.

Title: Progenitor cell markers predict outcome of patients with Hepatocellular Carcinoma beyond Milan criteria undergoing liver transplantation

Authors: Oriana Miltiadous¹, Daniela Sia^{1,2}, Yujin Hoshida¹, Maria Isabel Fiel¹, Andrew N. Harrington¹, Swan N. Thung¹, Poh Seng Tan^{1,3}, Hui Dong⁴, Kate Revill¹, Charissa Y Chang¹, Sasan Roayaie⁵, Thomas J Byrne⁶, Vincenzo Mazzaferro², Jorge Rakela⁶, Sander Florman¹, Myron Schwartz¹, and Josep M Llovet^{1,7,8}.

Affiliation:

¹ Mount Sinai Liver Cancer Program (Division of Liver Diseases, Department of Medicine, Tisch Cancer Institute, Department of Pathology, Recanati Miller Transplantation Institute, Department of Surgical Oncology), Icahn School of Medicine at Mount Sinai, New York, USA.

² Gastrointestinal Surgery and Liver Transplantation Unit, Department of Surgery, National Cancer Institute, Milan, Italy.

³ Division of Gastroenterology and Hepatology, University Medicine Cluster, National University Health System, Singapore

⁴ Eastern Hepatobiliary Surgery Hospital, Second Military Medical University Shanghai, China

⁵ Liver Cancer Program, Hofstra-North Shore LIJ School of Medicine, Lenox Hill Hospital, New York, USA

⁶ Division of Hepatology, Mayo Clinic, Phoenix, AZ, USA.

⁷ Liver Cancer Translational Research Laboratory, Barcelona - Clínic Liver Cancer Group (Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Liver Unit. Hospital Clínic, Universitat de Barcelona, Catalonia, Spain.

⁸ Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain.

Address for correspondence:

Josep M Llovet, MD
Mount Sinai Liver Cancer Program
Division of Liver Diseases,
Tisch Cancer Institute
Icahn School of Medicine at Mount Sinai.
Madison Ave 1425. 11F-70. Box 1123
New York, NY10029. USA
Phone: 212-6599503
FAX: 212-849-2574
E-mail: Josep.Llovet@mssm.edu

Electronic word count

Characters count Title: 121

Words count Abstract: 251

Words count includes Abstract, Introduction, Materials and Methods, Results,
Discussion, References, Tables and Figure Legends: 5985
References: 45

Number of Figures and Tables: 3 Figures and 5 Tables

Supplementary Material: 6 Figures and 6 Tables

List of abbreviations

HCC: Hepatocellular carcinoma
LT: Liver transplantation

1 CK19: Cytokeratin protein 19

2
3 HR: Hazard Ratio:

4
5 EpCAM: Epithelial cell adhesion molecule

6
7
8 OS: Overall survival

9
10 miVI: microvascular invasion

11
12 TGFb: Transforming growth factor beta

13
14 CT: Computed Tomography

15
16 MRI: Magnetic Resonance Imaging

17
18 AFP: Alpha-fetoprotein

19
20 IHC, immunohistochemical

21
22 RPS6: ribosomal protein S6

23
24 FDR: False discovery rate

25
26 H&E: Hematoxylin and eosin

27
28 PPV: Positive predictive value

29
30 NPV: Negative predictive value

31
32 HCV: Hepatitis C virus

33
34 HBV: Hepatitis B Virus

35
36 VIF: Variance Inflation Factor

37
38
39
40
41
42
43
44
45
46
47 **Keywords:** gene expression, prognosis, stem cell, gene signature, survival

48
49
50
51
52 **GEO Accession number:** GSE62743

53
54
55
56
57 **Conflict of Interest:** the authors indicate no potential conflicts of interest.

Financial Support:

Josep M. Llovet was supported by grants from The European Commission Framework Programme 7 (HEPTROMIC, Proposal No: 259744), the Samuel Waxman Cancer Research Foundation, the Spanish National Health Institute (SAF2013-41027), and the Asociación Española Contra el Cáncer (AECC). Daniela Sia and Vincenzo Mazzaferro are supported by the Italian Association for Cancer Research and the Italian National Ministry of Health. Yujin Hoshida is supported by grants from the U.S. National Institute of Diabetes and Digestive and Kidney Diseases (R01 DK099558), the European Commission Framework Programme 7 (HEPTROMIC, Proposal No: 259744).

Author's contributions

Conceived and designed the experiments: MS, JML, and JR

Performed the experiments: OM, DS, YH, ANH, PST, HD, and KR

Analyzed the data: OM, DS, YH, MIF, SNT, PST, CYC, JML

Contributed reagents/materials/analysis tools: OM, YH, ANH, PST, CYC, SR, TJB, VM, JR, SF, MS

Wrote the manuscript: OM, DS, YH, MS, JML

Clinical data: OM, ANH, CHC, SR, VM, JR, SF, MS, JML

Obtained funding: JR, SF, MS, JML

Critical review of the manuscript: OM, DS, YH, MIF, ANH, SNT, PST, KR, SR, TJB, VM, JR, SF, MS and JML

ABSTRACT

Background and Aims: In patients with hepatocellular carcinoma (HCC), liver transplantation (LT) is an excellent therapy if tumor characteristics are within the Milan criteria. We aimed to define genomic features enabling to identify HCC patients beyond Milan criteria who have acceptable transplant outcomes.

Methods: Among 770 consecutive HCC patients transplanted between 1990 and 2013, 132 had tumors exceeding Milan criteria on pathology and were enrolled in the study; 44% of the patients satisfied the 'up-to-7 rule' [7=sum of the size of the largest tumor and the number of tumors]. Explant tumors were assessed for genomic signatures and immunohistochemical markers associated with poor outcome.

Results: At a median follow-up of 88 months, 64 patients had died and 45 recurred; the 5-year overall survival (OS) and recurrence rates were 57% and 35%, respectively. Cytokeratin 19 (CK19) gene signature was independently associated with recurrence [Hazard ratio (HR)=2.95, $p<0.001$], along with tumor size (HR=3.37, $p=0.023$) and presence of satellites (HR=2.98, $p=0.001$). S2 subclass signature was independently associated with poor OS (HR=3.18, $p=0.001$), along with tumor size (HR=5.06, $p<0.001$) and up-to-7 rule (HR=2.50, $p=0.002$). Using the presence of progenitor cells markers (either CK19 or S2 signatures) patients were classified into poor-prognosis ($n=58$; 5-year recurrence 53%, survival 45%) and good-prognosis ($n=74$; 5-year recurrence 19%, survival 67%) (HR=3.16, $p<0.001$ for recurrence, and HR=1.72, $p=0.04$ for OS).

Conclusions:

HCC patients transplanted beyond Milan criteria without gene signatures of progenitor markers (CK19 and S2) achieved survival rates similar as those within Milan criteria.

Once prospectively validated, these markers may support a limited expansion of LT indications.

INTRODUCTION

Liver transplantation (LT) is an effective treatment option for hepatocellular carcinoma (HCC) when disease is defined by the widely accepted Milan criteria [1, 2]. Transplantation for patients within the Milan criteria generally yields a 5-year overall survival (OS) of 70% and a recurrence rate less than 15% [1-3]. Several efforts have been made to expand the criteria based on tumor size and number [4-8]. Although downstaging of tumors beyond Milan criteria is accepted by some UNOS regions, it has not yet been adopted by international consensus guidelines of LT [9] or guidelines of management of HCC [2]. Mazzaferro et al. proposed the up-to-7 rule [the sum of the number of tumor nodule(s) and the maximum diameter of the nodule(s) must not exceed the value of 7] which, in the absence of microvascular invasion (miVI) results in 5-year OS above 70% [8]. This study along with previously published data confirmed miVI to be a key predictor of recurrence in patients with HCC [10]. However, this information cannot be used in pre-transplant decision-making as an indication of transplantation since miVI is only diagnosed based on post-surgical histological assessment. This highlights the limitations of the current image-based prognostic algorithm for selecting HCC patients for LT.

Genome-wide transcriptome profiling has identified several key deregulated genes, molecular pathways and signatures associated with disease progression and prognosis in HCC [11, 12]. Activation of specific molecular pathways such as transforming growth factor-beta (TGFB) as well as presence of progenitor cell markers such as cytokeratin 19 (CK19) and epithelial cell adhesion molecule (EPCAM) have been associated with more aggressive biological tumor characteristics and rapid disease spread [13-35].

1 In a previous study, we identified gene signatures that significantly improved prediction
2 of HCC recurrence after surgical resection [21]. Here we sought to define if gene
3 signatures are also able to identify patients with HCC beyond Milan criteria who
4 nevertheless may have acceptable outcomes with LT. In addition, we are providing the
5 transcriptomic landscape of patients at more advanced states of the disease
6 compared with those undergoing resection or transplantation according to Milan
7 criteria, as per AASLD/EASL guidelines [2, 36].
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

MATERIALS AND METHODS

Patient cohorts and tissue samples

A total of 770 patients were transplanted for HCC between 1990 and 2013 at Mount Sinai Hospital, New York (n=590) and Mayo Clinic, Scottsdale-Arizona (n=180). Among these, we selected 132 patients (Mount Sinai: n=94, Mayo: n=38) with HCC beyond Milan criteria by pathological assessment, for whom base-line and follow-up clinical information and archived fixed tissue of viable tumor were available (**Figure 1**). The study protocol was approved by the respective Institutional Review Boards. See Supplementary Methods.

Genome-wide transcriptome profiling

Total RNA was subjected to transcriptome profiling using Whole-Genome DASL-HT (Illumina). See Supplementary Methods. GEO accession number: GSE62743.

Immunohistochemistry

Immunohistochemistry was performed for the evaluation of nuclear β -catenin as a surrogate for Wnt pathway activation; phosphorylation of ribosomal protein S6 (RPS6) as the major downstream effector of activated AKT-mTOR pathway; CK19 and EPCAM as surrogate of progenitor cell origin. See Supplementary Methods.

Bioinformatics and statistical analysis

Presence of previously reported HCC gene signatures (**Supplementary Table 1**) was evaluated (**Figure 2**). See Supplementary Methods.

RESULTS

Patient characteristics

Clinico-pathological data and outcomes of the patients are summarized in **Table 1**. Most of the patients were male (80%) and hepatitis C virus (HCV)-infected (71%). The median size of the largest tumor was 4 cm (range 1-9 cm). Tumors exceeded Milan criteria due to single tumor larger than 5 cm (5%), 2 to 3 tumors with the largest tumor larger than 3 cm (39%), or more than 3 tumors within the liver (57%). Satellite lesions (defined as HCC nodules < 2 cm in diameter within 2 cm from main tumor [37]) were present in 18% and miVI was present in 84% of the patients. Serum alpha fetoprotein (AFP) was > 200 mg/dL in 17% of cases. At the end of follow-up, 64 patients had died, OS at 5- and 10-years was 57% and 42%, respectively. HCC recurrence developed in 45 patients, 5 and 10-year cumulative recurrence rates were 35% and 44%, respectively (**Figure 3**).

Clinical variables identified as predictors of recurrence on multivariate analysis included tumor diameter >5 cm [$p=0.023$, Hazard Ratio (HR) = 3.37] and the presence of satellites ($p<0.001$, HR=2.98). Tumor diameter >5 cm ($p<0.001$, HR=5.06) and outside up-to-7 rule (HR=2.50, $p=0.002$) predicted poor survival (**Tables 2 and 3**).

Prevalence of prognostic liver cancer gene signatures

Among the 22 liver cancer gene signatures tested, 16 were able to sort patients into good and poor prognosis groups with a FDR<0.05 (**Figure 2A**). The human [21] and rat [28] CK19 gene signatures were present in 40% and 42% of patients, respectively. Signatures associated with poor clinical outcome and/or more aggressive biological features were generally present in the same set of tumors. At least one poor-prognosis signature was present in 84 patients (64%), this being a higher rate than we have

previously reported in patients treated with resection [21]. Twenty-five tumors (19%) harbored more than 10 poor prognosis/aggressiveness gene signatures. VI signature was positive in 22% of cases, 93% of which were positive for histological miVI. The majority of tumors lacking poor prognosis/aggressive signatures were enriched by the S3 subclass signature (39/48, 81%), which is associated with smaller, higher proportion of well-differentiated tumors [13].

The gene signatures analyzed were consistent with 3 distinct clusters or major molecular classes of HCC (**Figure 2B**): (1) *Proliferation and cell cycle activation* (Cluster A [16], Cholangiocarcinoma [29], Proliferation [20], Recurrence [24], S1 [13], TGFb [23], MET [18], and G3 [19] signatures), 2) *Proliferation-progenitor cell origin* (Hepatoblastoma C2 [26], Class S2 [13], EpCAM [25] and VI [22] signatures), which includes a subclass of CK19 positive tumors (CK19 human [21] and rat [28] signatures), and 3) *non-proliferative class*, enriched by tumors positive for CTNNB1[20] or class S3 [13, 33]. Interestingly, there was no association between the underlying etiology [HCV, Hepatitis B Virus (HBV) or alcohol] and the expression of signatures.

Comparing the classification of signatures between this cohort and previously published results from resection cases [21], we noticed a higher frequency of tumors positive for ≥ 6 signatures of poor prognosis (32.6% vs 25%, respectively). Similarly, our cohort presented a higher prevalence of CK19 human signatures (41% vs 34%, respectively), suggesting that it includes more advanced tumors compared with our previous report on resected patients [21].

Immunohistochemical (IHC) analysis

CK19 immunostaining was positive in 6 tumors (5%), and was significantly associated with the presence of the human ($p=0.03$) and rat ($p=0.04$) CK19 gene signatures, as well as VI signature [22] ($p<0.01$) (**Supplementary Table 2**). EPCAM immunostaining was positive in 20 tumors (15%), and was significantly associated with EPCAM signature ($p=0.01$), human and rat CK19 signatures ($p=0.01$ and $p=0.02$, respectively), Proliferation subclass signature ($p=0.01$), and VI signature ($p=0.01$) (**Supplementary Table 2**). Nuclear staining of β -catenin was observed in 27 tumors (21%), and was positively associated with CTNNB1 class signature ($p=0.01$) (**Supplementary Figure 1, Supplementary Table 2**). In addition, 12 samples (9%) had overexpression of cytoplasmic β catenin with one third of these in CTNNB1 class (4/12, 33%). Phospho-RPS6 staining was positive in 75 tumors (57%) with no obvious correlation with any of the evaluated gene signatures.

Prognostic relevance of liver cancer gene signatures in LT treated patients beyond Milan

Prognostic gene signatures in HCC were initially developed in patients undergoing resection, a treatment indicated for early stages of the disease. Univariate Cox regression analysis demonstrated that VI [22], G3 subclass [19], and S2 subclass [13] signatures were associated with overall survival, along with patient's age, tumor size, and up-to-7 rule (**Table 2**). In the multivariate analysis only S2 signature [13] ($HR=3.18$, $p=0.001$), tumor diameter >5 cm ($HR=5.06$, $p<0.001$), and outside up-to-7 rule ($HR=2.50$, $p=0.002$) were independent predictors of survival. VI [22], G3 [19], Proliferation [20], CK19 human [21], CK19 rat [28], Cholangiocarcinoma [29], Cluster A [16] and S3 [13] gene signatures were associated with HCC recurrence in univariate

analysis along with tumor size, AFP and satellites from clinical variables as well as RPS6 from IHC (**Table 3**). In multivariate analysis, CK19 human [21] (HR=2.91, p=0.001), tumor diameter > 5 cm (HR=3.37, p=0.023) and satellites (HR=2.98, p<0.001) were independent predictors of recurrence.

Of note, histological miVI predicted neither survival nor recurrence, presumably because in this cohort it was present in the large majority of cases (112 patients, 84.8%). Also, IHC positivity for reported progenitor cell markers also failed to improve outcome prediction.

Combination of progenitor cell signatures (CK19 and S2) discriminates prognostic subgroups in transplant patients

Multivariate analyses demonstrated prognostic significance of two progenitor-related gene signatures, CK19 [21] and S2 [13]. Furthermore, S2 subclass signature was also able to predict significant shorter survival post-recurrence in a subgroup analysis, suggesting that this signature might be able to capture more aggressive tumors with high risk of recurrence (p<0.001, HR= 5.11, **Supplementary Figure 2**). We therefore explored whether combining these signatures could further improve our ability to predict outcome. At least one of these two signatures was present in 58 patients (44%), and having one or both signatures independently predicted both OS (HR=1.72, p=0.036) and recurrence (HR=3.16, p=0.001) (**Tables 4 and 5**): patients with one or both signatures had 45% 5-year OS and 53% recurrence vs 67% 5-year OS and 19% recurrence when neither signature was present (**Figure 3C-D**). The presence of any of the two signatures also predicted recurrence when the analysis was limited to patients beyond Milan criteria based on preoperative imaging (n=94) (**Supplementary Figure 3, Supplementary Table 3**). Overall, 5-year recurrence rate was of 35.7% and was

significantly lower in patients CK19/S2 rule negative (20.8% vs 52.8%, $p=0.003$).

Furthermore, the CK19/S2 rule, when applied to patients classified according to BCLC stage, significantly improved the prognostic prediction by stratifying patients in high and low risk for both recurrence and survival (**Supplementary Figures 4 and 5**).

Discussion

Almost 20 years after the seminal observation by Mazzaferro et al [1], we continue to rely on size and number of tumors for selecting patients with HCC for LT. Since then, the concept of a modest expansion beyond the Milan criteria has been proposed without gaining global acceptance [4, 38-39]. Most of the studies performed were retrospective, used pathological data and only included few patients with HCC beyond Milan criteria, thus leading to unrealistic good results because the outcome was “diluted” among a much larger pool of patients within Milan criteria. The largest series ever reported exploring expansion of Milan criteria led to the pathological up-to-7 rule where outcomes were significantly better for patients within this rule, as long as vascular invasion is not present [8]. After all, tumor size and number remain surrogates for the biological aggressiveness of HCC; henceforth more refined markers are needed.

While there have been multiple publications linking various genetic profiles of HCC with outcomes after hepatic resection, tumor biology in transplanted patients has not been assessed widely thus far. Most studies have relied on degree of HCC differentiation on biopsy, AFP and response to bridge therapy while on the waiting list [40-42]. Using allelic imbalance, we described that genomic instability of certain microsatellites selected for proximity to known oncogenes can help select patients beyond Milan criteria who are likely to have favorable outcomes after transplantation [43]. Because gene expression profiling could be eventually performed on needle liver biopsies in a pre-transplant setting [44], the positivity for each of the signatures can assist in the decision making of selecting patients for transplantation.

In our current study, we explored the tumor gene expression profiles for the presence of 22 previously reported prognostic signatures –reflecting HCC molecular

subclasses or activation of signaling pathways- in 132 patients transplanted with HCC beyond Milan criteria and key immunohistochemical staining and correlated the results with clinical variables and outcome. Sixteen signatures demonstrated association with recurrence and/or survival; among these the CK19 and the S2 signatures, both denoting progenitor cell features, were independent predictors of outcome when included in models incorporating clinical variables. As shown in **Figure 2**, both signatures cluster together along with others from hepatoblastoma, which overall reflect progenitor cell origin. In fact, CK19 is a molecular marker of progenitor cells, whereas S2 subclass [13] has been linked with high levels of expression of AFP even at early stages of the disease, and activation of mTOR and IGF signaling [21, 35]. We also found enrichment for Met and TGFb signaling in these patients. In our study, one and/or both signatures were present in 44% of cases and the overall 5 yr-survival and recurrence rate (53% and 45%, respectively) was discouraging. Conversely, in the 56% of patients harboring neither signature, OS was 67% and recurrence was 19% at 5 years, similar to the results of LT for HCC when conventional Milan criteria are applied. The capacity of our tool to rule out tumors with “poor biology” is directly reflected in the acceptable low recurrence rate.

In order to be used prospectively, the CK19/S2 gene signature needs to be validated when Milan criteria are defined on pre-operative imaging. Indeed, in the subgroup analysis of pre-operatively defined cases beyond Milan criteria, the CK19/S2 signature remained independently associated with recurrence. However, the signature no longer retained significance on multivariate analysis for overall survival, likely due to the smaller sample size. Overall, these findings support the possibility that genomic analysis of HCC performed prior to LT may enable expansion of the current imaging-

1 based selection criteria to include patients beyond Milan Criteria but with favorable
2 genomic profiles, without significantly impacting clinical outcome.

3
4
5 Another interesting finding of our study was the relative genomic disparity observed in
6
7 HCC cases treated with LT versus those undergoing resection. In this series of
8
9 patients with HCC beyond Milan Criteria we observed a higher frequency of tumors
10
11 demonstrating ≥ 6 signatures than we did in the study of patients undergoing liver
12
13 resection [21]. This may be explained, to some degree, by the more advanced clinical
14
15 stage of cancers that received LT beyond Milan criteria versus those undergoing
16
17 resection. While resection is limited to single tumors, nearly 95% of the patients in the
18
19 current study were transplanted for multiple HCC.
20
21
22
23

24
25 We intend to validate these observations prospectively, but in order to establish
26
27 the absence of the CK19 and S2 progenitor cell signatures as a basis for safe
28
29 expansion of the indication for LT for patients with HCC, a number of barriers must be
30
31 overcome. First, the study will have to be limited to patients identified pre-LT as being
32
33 beyond Milan Criteria in order to be clinically relevant. The genomic heterogeneity of
34
35 tumors is widely recognized, all the more so when the tumor is large or multinodular as
36
37 in patients beyond Milan Criteria. While we have shown in a limited study that genomic
38
39 heterogeneity in HCC rarely results in assignment of samples from the same tumor to
40
41 different genomic classes [21], the degree to which a pre-LT biopsy accurately reflects
42
43 the genomic landscape of the tumor remains unclear. Molecular heterogeneity
44
45 between and within tumors can pose a risk for miss-classifications when selecting the
46
47 area to biopsy prior LT. However, with the advancements of imaging techniques, a
48
49 significant correlation has been shown between tumor enhancement and histological
50
51 differentiation which could greatly assist in more accurate profiling [45].
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 On the other side, the impact on gene expression of pre-LT nonsurgical tumor
2 treatment, that is commonly performed to down-stage HCC and forestalls progression
3 while awaiting transplant, remains undefined. Furthermore, with long pre-LT waiting
4 times before transplantation and the potential drop out of some patients, the stability of
5 tumor gene expression profiles over time must be considered. Thus, the extent to
6 which a biopsy done today reflects the nature of the tumor later on and the frequency
7 with which biopsy will need to be repeated, need to be determined.

8
9
10
11
12
13
14
15
16
17 Despite its shortcomings, our observations provide initial insights into the
18 understanding of tumor biology and clinical outcomes beyond size and number of
19 HCC nodules. Those patients with tumors with “good biology” as defined by absence
20 of signatures with progenitor cell traits (CK19/S2) represent half of the patients
21 transplanted beyond Milan criteria in 2 institutions in the United States. Our seminal
22 study provides the rationale for expanding the conventional criteria based upon
23 genomic data, since outcomes are competitive with those achieved by HCC patients
24 within Milan criteria. It is a novel, genomic, complementary tool to refine previously
25 reported proposals, such as Up-to-7 rule [8]. Certainly, these results need to be
26 validated using a separate external cohort of patients and then examined
27 prospectively before they can be adopted by clinical guidelines.

REFERENCES

- [1] Mazzaferro V, Regalia E, Doci R, Andreola S, Pulvirenti A, Bozzetti F et al. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med* 1996;334:693-700.
- [2] Llovet JM, Ducreux M, Lencioni R, Di Bisceglie AM, Galle PR, Dufour JF et al. J EASL-EORTC clinical practice guidelines: management of hepatocellular carcinoma. *Hepatology* 2012;56:908-43.
- [3] Llovet JM, Fuster J, Bruix J. Intention-to-treat analysis of surgical treatment for early hepatocellular carcinoma: resection versus transplantation. *Hepatology* 1999;30:1434-40.
- [4] Yao FY, Ferrell L, Bass NM, Watson JJ, Bacchetti P, Venook A, et al. Liver transplantation for hepatocellular carcinoma: expansion of the tumor size limits does not adversely impact survival. *Hepatology* 2001;33:1394-403.
- [5] Roayaie S, Frischer JS, Emre SH, Fishbein TM, Sheiner PA, Sung M et al. Long-term results with multimodal adjuvant therapy and liver transplantation for the treatment of hepatocellular carcinoma larger than 5 centimeters. *Ann Surg.* 2002;235:533–539.
- [6] Lee SG, Hwang S, Moon DB, Ahn CS, Kim KH, Sung KB et al. Expanded Indication Criteria of Living Donor Liver transplantation for Hepatocellular Carcinoma at One Large-Volume Center. *Liver Transpl.* 2008;14:935-45.
- [7] Yao FY. Liver transplantation for hepatocellular carcinoma: Beyond the Milan criteria. *Am J Transplant* 2008;8:1982-9.
- [8] Mazzaferro V, Llovet JM, Miceli R, Bhoori S, Schiavo M, Mariani L et al. Predicting survival after liver transplantation in patients with hepatocellular carcinoma beyond the Milan criteria: a retrospective, exploratory analysis. *Lancet Oncol.* 2009;10:35-43.

- 1 [9] Clavien PA, Lesurtel M, Bossuyt PM, Gores GJ, Langer B, Perrier A; OLT for HCC
2
3 Consensus Group. Recommendations for liver transplantation for hepatocellular
4
5 carcinoma: an international consensus conference report. *Lancet Oncol* 2012;13:e11-
6
7 22.
8
9
- 10 [10] Jonas S, Bechstein WO, Steinmüller T, Herrmann M, Radke C, Berg T, et al.
11
12 Vascular invasion and histopathologic grading determine outcome after liver
13
14 transplantation for hepatocellular carcinoma in cirrhosis. *Hepatology* 2001;33:1080-6.
15
16
- 17 [11] Villanueva A, Hernandez-Gea V, Llovet JM. Medical therapies for hepatocellular
18
19 carcinoma: a critical view of the evidence. *Nature Rev Gastroenterol Hepatol*
20
21 2013;10:34-42.
22
23
- 24 [12] Sawyer CL. The cancer biomarker problem, *Nature* 2008;452:548-52.
25
26
- 27 [13] Hoshida Y, Nijman SM, Kobayashi M, Chan JA, Brunet JP, Chiang DY et al.
28
29 Integrative Transcriptome Analysis Reveals Common Molecular Subclasses of Human
30
31 Hepatocellular Carcinoma. *Cancer Res* 2009;69:7385-92.
32
33
- 34 [14] Ye QH, Qin LX, Forgues M, He P, Kim JW, Peng AC et al. Predicting hepatitis B
35
36 virus-positive metastatic hepatocellular carcinomas using gene expression profiling
37
38 and supervised machine learning. *Nat Med* 2003;9:416-23.
39
40
- 41 [15] Iizuka N, Oka M, Yamada-Okabe H, Nishida M, Maeda Y, Mori N et al.
42
43 Oligonucleotide microarray for prediction of early intrahepatic recurrence of
44
45 hepatocellular carcinoma after curative resection. *Lancet* 2003;361:923-9.
46
47
- 48 [16] Lee JS, Chu IS, Heo J, Calvisi DF, Sun Z, Roskams T et al. Classification and
49
50 prediction of survival in hepatocellular carcinoma by gene expression profiling.
51
52 *Hepatology* 2004;40:667-76.
53
54
55
56
57
58
59
60
61
62
63
64
65

- [17] Kurokawa Y, Matoba R, Takemasa I, Nagano H, Dono K, Nakamori S et al. Molecular-based prediction of early recurrence in hepatocellular carcinoma. *J Hepatol*. 2004;41:284-91.
- [18] Kaposi-Novak P, Lee JS, Gómez-Quiroz L, Coulouarn C, Factor VM, Thorgeirsson SS. Met-regulated expression signature defines a subset of human hepatocellular carcinomas with poor prognosis and aggressive phenotype. *J Clin Invest*, 2006;116:1582-95.
- [19] Boyault S, Rickman DS, de Reyniès A, Balabaud C, Rebouissou S, Jeannot E, et al. Transcriptome classification of HCC is related to gene alterations and to new therapeutic targets. *Hepatology* 2007;45:42-52.
- [20] Chiang DY, Villanueva A, Hoshida Y, Peix J, Newell P, Minguez B, et al. Focal gains of VEGFA and molecular classification of hepatocellular carcinoma. *Cancer Res*. 2008;68:6779-88.
- [21] Villanueva A, Hoshida Y, Battiston C, Tovar V, Sia D, Alsinet C, et al. Combining clinical, pathology, and gene expression data to predict recurrence of hepatocellular carcinoma. *Gastroenterology*, 2011;140:1501-12.
- [22] Mínguez B, Hoshida Y, Villanueva A, Toffanin S, Cabellos L, Thung S et al. Gene-expression signature of vascular invasion in hepatocellular carcinoma. *J Hepatol*. 2011;55:1325-31.
- [23] Coulouarn C, Factor VM, Thorgeirsson SS. Transforming growth factor-beta gene expression signature in mouse hepatocytes predicts clinical outcome in human cancer. *Hepatology* 2008;47:2059-67.
- [24] Woo HG, Park ES, Cheon JH, Kim JH, Lee JS, Park BJ, et al. Gene expression-based recurrence prediction of hepatitis B virus-related human hepatocellular carcinoma. *Clin Cancer Res* 2008;14:2056-64.

- [25] Yamashita T, Forgues M, Wang W, Kim JW, Ye Q, Jia H, et al. EpCAM and alpha-fetoprotein expression defines novel prognostic subtypes of hepatocellular carcinoma. *Cancer Res* 2008;68:1451-61.
- [26] Cairo S, Armengol C, De Reyniès A, Wei Y, Thomas E, Renard CA et al. Hepatic stem-like phenotype and interplay of Wnt/beta-catenin and Myc signaling in aggressive childhood liver cancer. *Cancer Cell*, 2008;14:471-84.
- [27] Yoshioka S, Takemasa I, Nagano H, Kittaka N, Noda T, Wada H et al. Molecular prediction of early recurrence after resection of hepatocellular carcinoma. *Eur J Cancer* 2009;45:881-9.
- [28] Andersen JB, Loi R, Perra A, Factor VM, Ledda-Columbano GM, Columbano A et al. Progenitor-derived hepatocellular carcinoma model in the rat. *Hepatology* 2010;51:1401-9.
- [29] Woo HG, Lee JH, Yoon JH, Kim CY, Lee HS, Jang JJ et al. Identification of a cholangiocarcinoma-like gene expression trait in hepatocellular carcinoma. *Cancer Res.* 2010;70:3034-41.
- [30] Roessler S, Long EL, Budhu A, Chen Y, Zhao X, Ji J et al. Integrative genomic identification of genes on 8p associated with hepatocellular carcinoma progression and patient survival. *Gastroenterology* 2012;142:957-966.
- [31] Nault JC, De Reyniès A, Villanueva A, Calderaro J, Rebouissou S, Couchy G et al. A hepatocellular carcinoma 5-gene score associated with survival of patients after liver resection. *Gastroenterology* 2013;145:176-87.
- [32] Komuta M, Spee B, Vander Borgh S, De Vos R, Verslype C, Aerts R et al. Clinicopathological study on cholangiolocellular carcinoma suggesting hepatic progenitor cell origin. *Hepatology* 2008;47:1544-56.

- [33] Lachenmayer A, Alsinet C, Savic R, Cabellos L, Toffanin S, Hoshida Y et al. Wnt-pathway activation in two molecular classes of hepatocellular carcinoma and experimental modulation by sorafenib. *Cancer Res* 2012;18:4997-5007
- [34] Bjornsti MA, Houghton PJ. The tor pathway: a target for cancer therapy. *Nat Rev Cancer*;4:335-48.
- [35] Villanueva A, Chiang DY, Newell P, Peix J, Thung S, Alsinet C et al. Pivotal role of mTOR signaling in hepatocellular carcinoma. *Gastroenterology* 2008;135:1972-83.
- [36] Bruix J, Sherman M; American Association for the Study of Liver Diseases. Management of hepatocellular carcinoma: an update. *Hepatology*. 2011;53:1020-2.
- [37] Roayaie S, Blume IN, Thung SN, Guido M, Fiel MI, Hiotis S et al. A system of classifying microvascular invasion to predict outcome after resection in patients with hepatocellular carcinoma. *Gastroenterology* 2009;137:850-5.
- [38] Kim PT, Onaca N, Chinnakotla S, Davis GL, Jennings LW, McKenna GJ, et al. Tumor biology and pre-transplant locoregional treatments determine outcomes in patients with T3 hepatocellular carcinoma undergoing liver transplantation. *Clin Transplant*. 2013;27:311-8
- [39] Cillo U, Vitale A, Grigoletto F, Gringeri E, D'Amico F, Valmasoni M, et al. Intention-to-treat analysis of liver transplantation in selected, aggressively treated HCC patients exceeding the Milan criteria. *Am J Transplant*. 2007;7:972-81.
- [40] DuBay D, Sandroussi C, Sandhu L, Cleary S, Guba M, Cattral MS, et al. Liver transplantation for advanced hepatocellular carcinoma using poor tumor differentiation on biopsy as an exclusion criterion. *Ann Surg*. 2011;253:166-72.
- [41] Toso C, Asthana S, Bigam DL, Shapiro AM, Kneteman NM. Reassessing selection criteria prior to liver transplantation for hepatocellular carcinoma utilizing the Scientific Registry of Transplant Recipients database. *Hepatology* 2009;49:832-8

[42] Yao FY, Kerlan RK Jr, Hirose R, Davern TJ, Bass NM, Feng S, et al. Excellent outcome following down-staging of hepatocellular carcinoma prior to liver transplantation: an intention-to-treat analysis. *Hepatology*. 2008;48:819-27.

[43] Schwartz M, Dvorchik I, Roayaie S, Fiel MI, Finkelstein S, Marsh JW, et al. Liver transplantation for hepatocellular carcinoma: extension of indications based on molecular markers. *J Hepatol*. 2008;49:581-8.

[44] Hoshida Y, Villanueva A, Kobayashi M, Peix J, Chiang DY, Camargo A et al. Gene expression in fixed tissues and outcome in hepatocellular carcinoma. *N Engl J Med* 2008;359:1995-2004.

[45] Banerjee S, Wang DS, Kim HJ, Sirlin CB, Chan MG, Korn RL et al. A computed tomography radiogenomic biomarker predicts microvascular invasion and clinical outcomes in hepatocellular carcinoma. *Hepatology* 2015 Apr 30.

Acknowledgements

We thank Laia Cabellos for technical assistance and Xiaochen Sun for bioinformatic support.

FIGURE LEGENDS

Figure 1: Flow chart of the study. The initial cohort included 770 patients that were consecutively transplanted for HCC at Mount Sinai Hospital, New York (n=590) and Mayo Clinic, Scottsdale Arizona (n=180). A total of 206 patients were beyond Milan criteria based on the explant pathology. Tumors with HCC-ICC features, diffused pattern, necrotic tissue, macro-vascular invasion or metastasis were excluded. From the remaining 141 cases, 6 tumors with poor-quality RNA were discarded. Total RNA from 135 tumors was subjected to transcriptome profiling. Three samples had poor quality profile and were excluded and eventually 132 tumors were tested for the presence of previously reported outcome-associated gene signatures.

Figure 2: Prediction of gene expression signatures. Only signatures that were able to assign patients into good and poor prognosis groups with $FDR > 0.05$ were included. **(A)** Each column represents different sample and each row different signature. Positivity of each signature is represented by red bars. Events (recurrence or 5-year death) are shown with black bars. **(B)** Visualization of cramer's V coefficient for the pair-wise comparison of gene signatures. The scale from blue to red represents the strength of correlation (red represents the highest correlation). The signatures are clustered according to their correlation.

Figure 3: Kaplan-Meier curves and estimates of overall survival and recurrence in patients beyond Milan undergoing liver transplantation (n=132). Top figures show overall survival and recurrence in this cohort. Patients positive for either of CK19 or S2 signatures (progenitor-cell signatures) were assigned in a poor-outcome group.

Lower panels show overall survival and recurrence in patients beyond Milan with or without CK19/S2 signatures. P-values are calculated based on the log-rank test.

Table 1: Description of the main clinical and pathological characteristics of HCC patients undergoing LT beyond Milan criteria according to center and overall cohort (n=132)

Variable	Overall cohort (n=132)	Mount Sinai (n=94)	Mayo Clinic (n=38)
	n (%)	n (%)	n (%)
n	132 (100)	94 (71.2)	38 (28.8)
Age>60	39 (29.5)	28 (29.8)	11 (28.9)
Gender (males)	105 (79.5)	75 (79.8)	30 (78.9)
Transplant type			
Cadaveric donor	106 (80.3)	77 (81.9)	29 (76.3)
Living Donor	23 (17.4)	15 (16.0)	8 (21.1)
Underlying etiology*			
HCV	94 (71.2)	69 (73.4)	25 (65.8)
HBV	19 (14.4)	17 (18.1)	2 (5.26)
Alcohol	24 (18.2)	15 (16.0)	9 (23.7)
Other	10 (7.6)	8 (6.4)	4 (10.5)
Larger tumor size			
≤3cm	37 (28.0)	27 (28.7)	10 (26.3)
3-5cm	63 (47.7)	45 (47.9)	18 (47.4)
≥5cm	32 (24.2)	22 (23.4)	10 (26.3)
Number of tumors			
1-2	39 (29.5)	25 (26.6)	14 (36.8)
3-4	40 (30.3)	29 (30.8)	11 (28.9)
>5	53 (40.2)	40 (42.6)	13 (34.2)
Reason for extended criteria			
single tumor >5cm tumor	6 (4.5)	4 (4.3)	2 (5.3)
2-3 tumors, larger >3cm	51 (38.6)	34 (36.2)	17 (44.7)
>3 tumors	74 (56.1)	55 (58.5)	19 (50)
Pathological markers			
Well Differentiated	15 (11.4)	8 (8.5)	7 (18.4)
Moderately Differentiated	52 (39.4)	32 (34.0)	20 (52.6)
Poorly Differentiated	65 (49.2)	54 (57.4)	11 (28.9)
Satellites	24 (18.2)	22 (23.4)	2 (5.26)
Microvascular invasion	112 (84.8)	90 (95.7)	22 (57.9)
Satisfying up-to-7 rule	56 (44.1)	37 (41.1)	19 (51.4)
Within Mil pre-op	33 (26)	11 (12.4)	22 (57.9)
Alpha-fetoprotein >100mg/dL	30 (22.7)	25 (26.6)	5 (13.2)
Alpha-fetoprotein >200mg/dL	23 (17.4)	19 (20.2)	4 (10.5)
Albumin (<3.5mg/dL)	87 (65.9)	68 (72.3)	19 (50)
Bilirubin (>1mg/dL)	100 (75.8)	77 (81.9)	23 (60.5)
Child Pugh			
Class A	53 (40.2)	29 (36.3)	24 (63.2)
Class B	36 (30.5)	28 (35.0)	8 (21)
Class C	29 (24.6)	23 (28.8)	6 (15.8)
BCLC*			
A	27 (23.9)	23 (29.9)	4 (11.1)
B	86 (76.1)	54 (70.1)	32 (88.9)
Events			
Deaths	64 (48.5)	51 (54.3)	13 (34.2)
Overall Recurrence	45 (34.1)	34 (36.2)	11 (28.9)
Early Recurrence (<2years)	30 (22.7)	24 (25.5)	6 (15.8)
Follow up (months)			
median[Q25-Q75]	88 (65-122)	98 (78-138)	66 (37-80)
5-year Overall survival (%)	57.0	54.5	64.7
5-year Recurrence (%)	34.5	35.7	31.7

*BCLC defined with pre-operative clinical and radiological variables. Because patients were treated with liver transplantation, Child C patients were not considered BCLC D and were relocated regardless of liver function.

Table 2: Univariate and Multivariate analysis of clinical, pathological and gene signatures for overall survival in patients beyond Milan criteria (n=132)

Gene expression signatures	Patients Pos Rest		Overall Survival, events n=64					
			Univariate analysis			Multivariate analysis		
			pval	HR	95% CI	pval	HR	95% CI
VI signature [22]	29	103	0.040	1.760	1.02-3.04	0.001	3.177	1.60-6.31
G3 signature [19]	40	92	0.040	1.705	1.02-2.86			
Proliferation [20]	39	93	0.250	1.356	0.80-2.29			
CK19 human [21]	53	79	0.054	1.611	0.99-2.63			
CK19 rat [28]	56	76	0.130	1.457	0.89-2.38			
MET signature [18]	34	98	0.458	0.800	0.44-1.45			
Hepatoblastoma_C2 [26]	6	126	0.086	2.369	0.86-6.56			
CTNNB1 [20]	30	102	0.208	1.435	0.81-2.53			
EpCAM [25]	26	106	0.411	1.281	0.71-2.32			
Cholangiocarcinoma [29]	44	88	0.471	1.209	0.72-2.03			
Recurrence [24]	41	91	0.713	1.104	0.65-1.88			
Cluster A [16]	45	87	0.369	1.264	0.76-2.11			
TGFb [23]	38	94	0.762	0.918	0.53-1.60			
S1 [13]	42	90	0.545	0.845	0.49-1.46			
S2 [13]	15	117	0.011	2.274	1.18-4.37			
S3 [13]	47	85	0.358	0.781	0.50-1.33			
IHC								
CK19	6	126	0.571	1.396	0.44-4.46			
EpCAM	20	112	0.096	0.497	0.21-1.15			
bcatenin	27	105	0.334	1.329	0.74-2.37			
RPS6	75	57	0.173	1.414	0.86-2.33			
Clinical Variables								
Male gender	105	27	0.441	0.775	0.40-1.49	<0.001	5.057	2.06-12.44
Age (>60)	39	93	0.031	1.730	1.04-2.87			
Etiology	94	38	0.168	1.512	0.84-2.74			
HCV	19	113	0.381	0.718	0.34-1.51			
HBV	24	108	0.363	1.316	0.73-2.39			
Alcohol	6	126	<0.001	4.289	1.83-10.07			
Single tumor >5cm	51	81	0.118	0.659	0.39-1.12			
2-3 tumors, largest >3cm	75	57	0.675	1.112	0.68-1.83			
No of tumors (>3)	23	109	0.087	1.650	0.92-2.95			
AFP (>200mg/dL)	112	20	0.051	2.631	0.96-7.24			
MiVI	24	108	0.056	1.724	0.98-3.04	0.002	2.499	1.42-4.42
Satellites	65	67	0.183	1.398	0.85-2.30			
Differentiation (Poor)	69	56	0.018	1.869	1.10-3.17			
Outside Up-to-7	32	95	0.531	0.818	0.44-1.54			
Within Milan in pre-op assessment	27	86	0.184	1.491	0.82-2.70			
BCLC A	53	65	0.322	0.758	0.44-1.32			
Child A								

Table 3: Univariate and Multivariate analysis of clinical, pathological and gene signatures for overall recurrence in patients beyond Milan criteria (n=132)

Gene expression signatures	Patients Pos Rest		Overall Recurrence, events n=45					
			Univariate analysis			Multivariate analysis		
			pval	HR	95% CI	pval	HR	95% CI
VI signature [22]	29	103	0.008	2.262	1.22-4.21	0.001	2.905	1.59-5.32
G3 signature [19]	40	92	0.038	1.861	1.02-3.39			
Proliferation [20]	39	93	0.004	2.297	1.28-4.14			
CK19 human [21]	53	79	<0.001	3.031	1.66-5.54			
CK19 rat [28]	56	76	0.001	2.676	1.46-4.90			
MET signature [18]	34	98	0.181	1.520	0.82-2.83			
Hepatoblastoma_C2 [26]	6	126	0.702	1.318	0.32-5.47			
CTNNB1 [20]	30	102	0.986	0.994	0.48-2.07			
EpCAM [25]	26	106	0.461	1.301	0.64-2.63			
Cholangiocarcinoma [29]	44	88	0.036	1.855	1.03-3.34			
Recurrence [24]	41	91	0.366	1.323	0.72-2.44			
Cluster A [16]	45	87	0.046	1.802	1.00-3.25			
TGFb [23]	38	94	0.917	1.035	0.54-1.97			
S1 [13]	42	90	0.529	1.215	0.66-2.24			
S2 [13]	15	117	0.132	1.842	0.82-4.14			
S3 [13]	47	85	0.012	0.408	0.20-0.85			
IHC								
CK19	6	126	0.321	1.610	0.38-6.85			
EpCAM	20	112	0.416	0.702	0.30-1.66			
bcatenin	27	105	0.381	1.353	0.68-2.68			
RPS6	75	57	0.025	2.024	1.07-3.82			
Clinical Variables								
Male gender	105	27	0.558	0.797	0.37-1.71	0.023	3.374	1.18-9.63
Age (>60)	39	93	0.782	1.095	0.57-2.09			
Etiology	HCV	94	0.254	1.499	0.74-3.03			
	HBV	19	0.204	0.521	0.19-1.46			
	Alcohol	24	0.679	1.170	0.56-2.42			
Single tumor >5cm	6	126	0.016	3.306	1.17-9.34			
2-3 tumors, largest >3cm	51	81	0.168	0.644	0.34-1.21			
No of tumors (>3)	75	57	0.580	1.183	0.65-2.15			
AFP (>200mg/dL)	23	109	0.003	2.579	1.35-4.93			
MiVI	112	20	0.181	1.982	0.71-5.54			
Satellites	24	108	<0.001	3.069	1.65-5.72	<0.001	2.978	1.60-5.55
Differentiation (Poor)	65	67	0.271	1.388	0.77-2.50			
Outside Up-to-7	69	56	0.241	1.444	0.78-2.67			
Within Milan in pre-op assessment	32	95	0.382	0.710	0.33-1.53			
BCLC A	27	86	0.962	1.018	0.48-2.15			
Child A	53	65	0.587	1.187	0.64-2.12			

Table 4: Univariate and Multivariate analysis of CK19/S2 score for overall survival in patients beyond Milan (n=132)

Gene expression Signatures	Patients Pos Rest		Overall Survival, events n=64					
			Univariate analysis			Multivariate analysis		
			pval	HR	95% CI	Pval	HR	95% CI
CK19 [21] and/or S2 pos [13]	58	74	0.030	1.17	1.04-2.79	0.036	1.719	1.034-2.85
IHC								
CK19	6	126	0.571	1.396	0.44-4.46			
EpCAM	20	112	0.096	0.497	0.21-1.15			
bcatenin	27	105	0.334	1.329	0.74-2.37			
RPS6	75	57	0.173	1.414	0.86-2.33			
Clinical Variables								
Male gender	105	27	0.441	0.775	0.40-1.49			
Age (>60)	39	93	0.031	1.730	1.04-2.87			
Etiology								
HCV	94	38	0.168	1.512	0.84-2.74			
HBV	19	113	0.381	0.718	0.34-1.51			
Alcohol	24	108	0.363	1.316	0.73-2.39			
Single tumor >5cm	6	126	<0.001	4.289	1.83-10.07	<0.001	5.510	2.26-13.45
Two-Three tumors, larger >3cm	51	81	0.118	0.659	0.39-1.12			
No of tumors (>3)	75	57	0.675	1.112	0.68-1.83			
AFP (>200mg/dL)	23	109	0.087	1.650	0.92-2.95			
MiVI	112	20	0.051	2.631	0.96-7.24			
Satellites	24	108	0.056	1.724	0.98-3.04			
Differentiation (Poor)	65	67	0.183	1.398	0.85-2.30			
Outside Up-to-7	69	56	0.018	1.869	1.10-3.17	0.007	2.113	1.23-3.64
Within Milan in pre-op assessment	32	95	0.531	0.818	0.44-1.54			
BCLC A	27	86	0.184	1.491	0.82-2.70			
Child A	53	65	0.322	0.758	0.44-1.32			

Table 5: Univariate and Multivariate analysis of CK19/S2 score for overall recurrence in patients beyond Milan (n=132)

Gene expression Signatures	Patients Pos Rest		Overall Recurrence, events n=45					
			Univariate analysis			Multivariate analysis		
			pval	HR	95% CI	pval	HR	95% CI
CK19 [21] and/or S2 pos [13]	58	74	<0.001	3.192	1.67-5.60	<0.001	3.162	1.70-5.89
IHC								
CK19	6	126	0.321	1.610	0.38-6.85			
EpCAM	20	112	0.416	0.702	0.30-1.66			
bcatenin	27	105	0.381	1.353	0.68-2.68			
RPS6	75	57	0.025	2.024	1.07-3.82			
Clinical Variables								
Male gender	105	27	0.558	0.797	0.37-1.71			
Age (>60)	39	93	0.782	1.095	0.57-2.09			
Etiology								
HCV	94	38	0.254	1.499	0.74-3.03			
HBV	19	113	0.204	0.521	0.19-1.46			
Alcohol	24	108	0.679	1.170	0.56-2.42			
Single tumor >5cm	6	126	0.016	3.306	1.17-9.34	0.019	3.520	1.23-10.06
Two-Three tumors, larger >3cm	51	81	0.168	0.644	0.34-1.21			
No of tumors (>3)	75	57	0.580	1.183	0.65-2.15			
AFP (>200mg/dL)	23	109	0.003	2.579	1.35-4.93			
MiVI	112	20	0.181	1.982	0.71-5.54			
Satellites	24	108	<0.001	3.069	1.65-5.72	<0.001	3.078	1.65-5.74
Differentiation (Poor)	65	67	0.271	1.388	0.77-2.50			
Outside Up-to-7	69	56	0.241	1.444	0.78-2.67			
Within Milan in pre-op assessment	32	95	0.382	0.710	0.33-1.53			
BCLC A	27	86	0.962	1.018	0.48-2.15			
Child A	53	65	0.587	1.187	0.64-2.12			

Fig. 1
Figure 1





Fig. 2
Figure 2

A.



B.





Figure 3


A.

Overall Survival




B.

Recurrence




C.

Overall Survival



D.

Recurrence



Supplementary material

[Click here to download Supplementary material: Supplementary_Miltiadous_Miltiadous_Llovet_July2015.docx](#)